

Symmetric exclusion as a random environment: Hydrodynamic limits

Luca Avena^a, Tertuliano Franco^b, Milton Jara^c and Florian Völlering^d

^a*Institut für Mathematik, Universität Zürich, Switzerland. E-mail: luca.avena@math.uzh.ch*

^b*Universidade Federal da Bahia, Salvador, Brazil. E-mail: tertu@impa.br*

^c*IMPA, Rio de Janeiro, Brazil. E-mail: mjara@impa.br*

^d*Mathematisch Instituut, Universiteit Leiden, The Netherlands. E-mail: fvolleri@math.leidenuniv.nl*

Received 10 July 2013; revised 25 November 2013; accepted 23 January 2014

Abstract. We consider a one-dimensional continuous time random walk with transition rates depending on an underlying autonomous simple symmetric exclusion process starting out of equilibrium. This model represents an example of a random walk in a slowly non-uniform mixing dynamic random environment. Under a proper space–time rescaling in which the exclusion is speeded up compared to the random walk, we prove a hydrodynamic limit theorem for the exclusion as seen by this walk and we derive an ODE describing the macroscopic evolution of the walk. The main difficulty is the proof of a replacement lemma for the exclusion as seen from the walk without explicit knowledge of its invariant measures. We further discuss how to obtain similar results for several variants of this model.

Résumé. Nous considérons une marche aléatoire unidimensionnelle à temps continu, avec des taux des sauts dépendants d'un processus d'exclusion autonome et hors équilibre. Ce modèle représente un exemple de marche aléatoire en milieu aléatoire dynamique, où le milieu n'a pas des bonnes propriétés de mélange. Sous la bonne échelle spatio-temporelle, où le processus d'exclusion est accéléré de plus en plus par rapport à la marche, nous démontrons un théorème de limite hydrodynamique pour le processus d'exclusion vu par la marche aléatoire, et nous dérivons une EDO qui décrit l'évolution macroscopique de la marche. La difficulté principale est la démonstration d'un lemme de remplacement pour le processus d'exclusion vu par la marche aléatoire, sans une connaissance explicite de ses mesures invariantes. Nous discutons comment obtenir des résultats similaires pour des variantes du modèle en question.

MSC: 82C05; 60K37

Keywords: Random walks in random environments; Macroscopic speed; Hydrodynamic limits; Particle systems; Exclusion process

1. Introduction

1.1. Model and motivation

Let $\Omega = \{0, 1\}^{\mathbb{Z}}$. Denote by $\eta = \{\eta(z); z \in \mathbb{Z}\}$ the elements of Ω . For $\eta \in \Omega$ and $z \in \mathbb{Z}$, define $\eta^{z, z+1} \in \Omega$ as

$$\eta^{z, z+1}(x) = \begin{cases} \eta(z+1), & x = z, \\ \eta(z), & x = z+1, \\ \eta(x), & x \neq z, z+1, \end{cases}$$

that is, $\eta^{z,z+1}$ is obtained from η by exchanging the occupation variables at z and $z+1$. Fix $\alpha, \beta \geq 0$ and assume that $\alpha + \beta > 0$. Let $\{(\eta_t, x_t); t \geq 0\}$ be the Markov process on the state space $\Omega \times \mathbb{Z}$ with generator given by

$$\begin{aligned} Lf(\eta, x) &= \sum_{z \in \mathbb{Z}} [f(\eta^{z,z+1}, x) - f(\eta, x)] + \sum_{y \in \{\pm 1\}} c_y(\eta(x)) [f(\eta, x+y) - f(\eta, x)] \\ &:= \sum_{z \in \mathbb{Z}} [f(\eta^{z,z+1}, x) - f(\eta, x)] + [\beta + (\alpha - \beta)\eta(x)] [f(\eta, x+1) - f(\eta, x)] \\ &\quad + [\alpha + (\beta - \alpha)\eta(x)] [f(\eta, x-1) - f(\eta, x)], \end{aligned} \tag{1.1}$$

for any local function $f: \Omega \times \mathbb{Z} \rightarrow \mathbb{R}$. We interpret the dynamics of the process $\{(\eta_t, x_t); t \geq 0\}$ as follows. Checking the action of L over functions f which do not depend on x , we see that $\{\eta_t; t \geq 0\}$ has a Markovian evolution, which corresponds to the well-known *simple symmetric exclusion process* on \mathbb{Z} . Conditioned on a realization of $\{\eta_t; t \geq 0\}$, the process $\{x_t; t \geq 0\}$ is a random walk that jumps to the left with rate β and to the right with rate α whenever there is a particle at the current position of the random walk (i.e., $\eta(x_t) = 1$). When there is no particle at the current position of the random walk (i.e., $\eta(x_t) = 0$), the rates are reversed: it jumps to the left with rate α and it jumps to the right with rate β . We say that the simple exclusion process $\{\eta_t; t \geq 0\}$ is a *dynamical random environment* and that $\{x_t; t \geq 0\}$ is a random walk in such a dynamical random environment. Note that the fact the random walk has a local drift $\alpha - \beta$ on occupied sites, and opposite drift $\beta - \alpha$ on vacant sites, creates trapping phenomena typical of random walks in random environments. This random walk has a tendency to spend a long time around the interface between regions with majority of particles and regions with majority of holes. As we will show, our proofs and results are still valid for more general choices of the local drifts of the process $\{x_t; t \geq 0\}$, see Remark 1.4 and Section 3.2 below.

Our main results are Theorem 1.2 and Theorem 1.3 below. Informally speaking, our results can be formulated as follows. From the hydrodynamic limit theory for the exclusion process it is known that when time is rescaled by N^2 and space is rescaled by $1/N$, then the sequence of rescaled exclusion processes converges to the heat equation. We will rescale the random walk driven by the exclusion process differently, namely by N in time and $1/N$ in space (see (1.6) for the precise definition). Then we consider the joint sequence of rescaled random walk and rescaled exclusion process and obtain an ODE governing the macroscopic evolution of the random walk limit, and a hydrodynamic limit theorem for the exclusion as seen by this walker.

In Section 3 we will discuss how to derive the same results for several variants of this model. In order to derive Theorem 1.3 we need to prove a so-called *replacement lemma*, Theorem 2.1, in the spirit of [10]. Unlike [10], we do not have explicit knowledge of the invariant measures for the particle system given by the environment as seen by the walker. The latter is a notoriously difficult problem in hydrodynamic limit theory. However, in our setting, the Bernoulli product measures (which are invariant for the exclusion process) turn out to be close to being invariant for the environment as seen by the walker under the mentioned rescaling. In fact, we show a crucial estimate in Lemma 2.2, which will allow to control the entropy between the distribution of the evolved environment as seen by the walker and the Bernoulli product measures. This estimate is enough to prove the mentioned *replacement lemma*, Theorem 2.1. See the paragraph below Theorem 2.1 for further explanations.

In recent years, there has been much interest in the study of random walks in random environments. See, e.g., [13,14] and [3,5,7] for recent results, overviews and references in *static* and *dynamic* environments, respectively. The aim is to understand the motion of a particle in a material presenting impurities which is of clear interest for applied purposes. Despite of the increasing literature on the subject, several basic questions are still open even in one dimension. The model we analyse in this paper has been introduced in [1] as a model of a random walk in a Markovian autonomous environment with *slow* and *non-uniform* mixing properties due to the fact that space–time correlations in the exclusion decay slowly and there is not a unique invariant measure. Almost no rigorous results are known in the latter setting since the general techniques and results in the literature are suitable only for *fast uniform* mixing types of environments.

From a phenomenological point of view, when considering *fast uniform* mixing type of environments (e.g., if $\{\eta_t; t \geq 0\}$ evolves according to an independent spin flips dynamics) trapping effects play a minor role. Indeed the asymptotic behavior of $\{x_t; t \geq 0\}$ in such a setting is qualitatively equivalent to the one of a homogeneous random walk, namely, ballisticity in the transient regime, diffusive scaling and exponential decay for large deviations. In [1], the authors considered this model characterized by (1.1) showing sub-exponential cost for sub-linear displacement of

$\{x_t; t \geq 0\}$. In other words, despite of the dynamics of the environment, this model shows some slow-down phenomena similar to a static random environment (see, e.g., [6] and [4]). When looking at fluctuations, as suggested by the numerics in [3], non-diffusive behavior is also expected. Therefore *slowly non-uniform* mixing environments behave very differently from *fast uniform* mixing ones.

Further rigorous results on this model have been derived in [2] where the authors considered the case where the process $\{x_t; t \geq 0\}$ has still two different drifts on occupied and vacant sites of the exclusion but both local drifts in the same direction. Under this latter assumption, traps do not play any role and standard diffusive scaling and exponential decay of deviations from the typical behavior have been proven by means of a delicate renewal construction.

Unfortunately, the type of rescaling (see (1.6)) required in our Theorem 1.2 does not imply the law of large numbers for the original random walk considered in [1,3]. In fact, to pass from a micro to a macroscopic scale, we need that the space–time rescaling of the two processes $\{x_t; t \geq 0\}$ and $\{\eta_t; t \geq 0\}$ are of the same order. In other words, under our space–time rescaling in (1.6), the exclusion jumps much faster than the walk preventing “strong trapping effects.” Nevertheless, on one hand, our results give a new contribution in the field of random walks in *slowly non-uniform* mixing dynamic random environments. On the other hand, they strengthen the connection between random walks in random environments and scaling limits of particle systems. In this respect, there is a wide literature on the problem of the tagged particle in conservative particle systems, see [12] for a review on this subject. The tagged particle can be also interpreted as a random walk in a random environment with two main differences with respect to our model: walker and environment are mutually interacting, and the invariant measures of the environment as seen by the walker are explicitly known. Our Theorem 1.3 is also of interest in itself within the hydrodynamic limit theory because of this lack of knowledge for the invariant measures of the environment from the point of the walker. In the next section we define this process before finally stating the results.

1.2. The environment as seen by the walker

Consider the process $\{\xi_t; t \geq 0\}$ with values in Ω , defined by $\xi_t(z) = \eta_t(z + x_t)$. The process $\{\xi_t; t \geq 0\}$ is called the *environment as seen by the random walk*. In other words, $\xi_t = \theta^{x_t} \eta$, where θ^z denotes the shift operator for $z \in \mathbb{Z}$. It turns out that the process $\{\xi_t; t \geq 0\}$ is a Markov process. Its generator is given by

$$\mathcal{L}f(\xi) = \sum_{z \in \mathbb{Z}} [f(\xi^{z, z+1}) - f(\xi)] + \sum_{y \in \{\pm 1\}} c_y(\xi(0)) [f(\theta^y \xi) - f(\xi)] \quad (1.2)$$

for any local function $f : \Omega \rightarrow \mathbb{R}$, with

$$c_{+1}(\xi(0)) = \beta + (\alpha - \beta)\xi(0) \quad \text{and} \quad c_{-1}(\xi(0)) = \alpha + (\beta - \alpha)\xi(0). \quad (1.3)$$

The dynamics of $\{\xi_t; t \geq 0\}$ is the following. Particles move according to a simple symmetric exclusion process. Superposed to this dynamics, the configuration of particles is shifted to the left or to the right with rates corresponding to the jumps of the random walk $\{x_t; t \geq 0\}$.

1.3. Main results: Scaling limits

In order to state our main results, we first introduce some notation and recall what is known in the literature as the hydrodynamic limit for the simple symmetric exclusion process. Let $u_0 : \mathbb{R} \rightarrow [0, 1]$ be a piecewise continuous function and let $n \in \mathbb{N}$ be a scaling parameter. We define a probability measure μ^n in Ω by

$$\mu^n(\eta(z_1) = 1, \dots, \eta(z_\ell) = 1) = \prod_{i=1}^{\ell} u_0(z_i/n) \quad (1.4)$$

for any set $\{z_1, \dots, z_\ell\} \subseteq \mathbb{Z}$. We call the sequence of measures $\{\mu^n\}_{n \in \mathbb{N}}$, the *Bernoulli product measures associated to the profile u_0* . Next, let us define the *empirical measure* $\{\pi_t^n; t \geq 0\}$ as the measure-valued process given by

$$\pi_t^n(dx) = \frac{1}{n} \sum_{z \in \mathbb{Z}} \eta_t^n(z) \delta_{z/n}(dx),$$

where $\delta_{z/n}(dx)$ denotes the Dirac measure at $\frac{z}{n} \in \mathbb{R}$, and $\{\eta_t^n; t \geq 0\}$ denotes the process $\{\eta_{tn^2}; t \geq 0\}$.

The following result is known in the literature as the *hydrodynamic limit* of the simple symmetric exclusion process, see, e.g., Theorem 2.1, Chapter 2 in [11].

Proposition 1.1 (Hydrodynamic equation for the simple exclusion). *Let $u_0 : \mathbb{R} \rightarrow [0, 1]$ be a piecewise continuous function and let $\{\mu^n\}_{n \in \mathbb{N}}$ be the sequence of Bernoulli product measures associated to the profile u_0 as in (1.4). Let $\{\eta_t^n; t \geq 0\}$ the process $\{\eta_{tn^2}; t \geq 0\}$ with initial distribution μ^n . For any $T \geq 0$, the sequence of measure-valued processes $\{\pi_t^n(dx); t \in [0, T]\}_{n \in \mathbb{N}}$ converges to $\{u(t, x) dx; t \in [0, T]\}$ in probability with respect to the J_1 -Skorohod topology of the space of càdlàg paths $\mathbb{D}([0, T]; \mathcal{M}_+(\mathbb{R}))$, where $\{u(t, x); t \geq 0, x \in \mathbb{R}\}$ is the solution of the Cauchy problem*

$$\begin{cases} \partial_t u(t, x) = (1/2)\partial_{xx}^2 u(t, x), & t \geq 0, x \in \mathbb{R}, \\ u(0, x) = u_0(x), & x \in \mathbb{R}. \end{cases} \tag{1.5}$$

Back to our model in (1.1), heuristically, it is easy to see that if the average density of particles is different from $\frac{1}{2}$, the random walk $\{x_t; t \geq 0\}$ moves ballistically, and therefore the diffusive scaling introduced above is not the right one for $\{x_t; t \geq 0\}$. A possible way to overcome this fact is to scale the exclusion process and the random walk in a different way. One way to do this is to define the rescaled process $\{\eta_t^n, x_t^n; t \geq 0\}$ as the Markov process generated by the operator

$$\begin{aligned} L_n f(\eta, x) &= n^2 \sum_{z \in \mathbb{Z}} [f(\eta^{z, z+1}, x) - f(\eta, x)] \\ &+ n \sum_{y \in \{\pm 1\}} c_y(\eta(x)) [f(\eta, x + y) - f(\eta, x)]. \end{aligned} \tag{1.6}$$

Under such a rescaling the exclusion jumps faster than the random walk but the motion of the random walk and of the particles in the exclusion occur at the same scale allowing for a macroscopic non-trivial description of their evolution. We are finally in shape to state the first result concerning $\{x_t^n; t \geq 0\}$.

Theorem 1.2 (Macroscopic evolution of the random walk). *Let $u_0 : \mathbb{R} \rightarrow [0, 1]$ be a piecewise continuous function and let $\{\mu^n\}_{n \in \mathbb{N}}$ be the sequence of Bernoulli product measures associated to the profile u_0 . Assume that there exists a constant $\rho \in (0, 1)$ such that*

$$\sup_n \frac{1}{n} \sum_{x \in \mathbb{Z}^d} \left| u_0\left(\frac{x}{n}\right) - \rho \right|^2 < +\infty. \tag{1.7}$$

Fix $x_0^n = 0$ and $T > 0$. Let $\{\eta_t^n; t \geq 0\}$ be the process $\{\eta_{tn^2}; t \geq 0\}$ with initial distribution μ^n . Then, for all $t \in [0, T]$:

$$\lim_{n \rightarrow \infty} \frac{x_t^n}{n} = f(t)$$

in probability, where $\{f(t); t \geq 0\}$ satisfies the ordinary differential equation

$$\begin{cases} f'(t) = (\beta - \alpha)(1 - 2u(t, f(t))), \\ f(0) = 0, \end{cases} \tag{1.8}$$

with u being the solution of (1.5).

Note that by assuming (1.7), we do not restrict ourselves to the case where the exclusion starts from equilibrium, namely, when the initial profile u_0 is constantly equal to some fix density ρ . When restricting to this case for the starting density profile, i.e., $u_0 \equiv \rho$, Theorem 1.2 reduces to the statement that the random walk observes an averaged homogeneous environment and travels with constant speed given by

$$f'(t) = (\beta - \alpha)(1 - 2\rho).$$

In other words, when $u_0 \equiv \rho$, it is like an homogeneous random walk jumping with probabilities $\beta + \rho(\alpha - \beta)$ and $\alpha + \rho(\beta - \alpha)$ to the right and to the left, respectively.

The proof of this theorem will be linked to the proof of the following theorem, which represents the hydrodynamic limit of the environment as seen by the random walk. Let $\{\xi_t^n; t \geq 0\}$ be the rescaled environment as seen by the walk given by $\xi_t^n(x) = \eta_t^n(x + x_t^n)$. Its generator \mathcal{L}_n is given by

$$\begin{aligned} \mathcal{L}_n f(\xi) &= n^2 \sum_{z \in \mathbb{Z}} [f(\xi^{z, z+1}) - f(\xi)] + n \sum_{y \in \{\pm 1\}} c_y(\xi(0)) [f(\theta^y \xi) - f(\xi)] \\ &=: n^2 \mathcal{L}^{\text{ex}} f(\xi) + n \mathcal{L}^{\text{rw}} f(\xi). \end{aligned} \tag{1.9}$$

Let $\{\hat{\pi}_t^n(dx); t \geq 0\}$ be the empirical measure associated to the process $\{\xi_t^n; t \geq 0\}$:

$$\hat{\pi}_t^n(dx) = \frac{1}{n} \sum_{z \in \mathbb{Z}} \xi_t^n(z) \delta_{z/n}(dx). \tag{1.10}$$

We have the following:

Theorem 1.3 (Hydrodynamic equation for the environment as seen by the walker). Fix $T \geq 0$, under the assumptions of Theorem 1.2, the sequence of measure-valued processes $\{\hat{\pi}_t^n(dx); t \in [0, T]\}_{n \in \mathbb{N}}$ converges to $\{\hat{u}(t, x) dx; t \in [0, T]\}$ in probability with respect to the J_1 -Skorohod topology of $\mathbb{D}([0, T]; \mathcal{M}_+(\mathbb{R}))$, where $\{\hat{u}(t, x); t \geq 0, x \in \mathbb{R}\}$ is the solution of the Cauchy problem

$$\begin{cases} \partial_t \hat{u}(t, x) = (1/2) \partial_{xx}^2 \hat{u}(t, x) + (\beta - \alpha)(1 - 2\hat{u}(t, 0)) \partial_x \hat{u}(t, x), & t \geq 0, x \in \mathbb{R}, \\ \hat{u}(0, x) = u_0(x), & x \in \mathbb{R}. \end{cases} \tag{1.11}$$

Theorem 1.3 says that the effect of the random walk on the macroscopic evolution of the process $\{\xi_t^n; t \geq 0\}$ results into a transport term. This term reflects the fact that microscopically every time the random walk jumps to the right, the exclusion gets shifted to the left, and vice versa. The speed of this transport mechanism is the speed of the walker that is dependent on $\hat{u}(t, 0) = u(t, f(t))$, which represents the density observed by $\{x_t^n; t \geq 0\}$.

Let us further stress that Theorem 1.3 has its own interest within hydrodynamic theory of particle systems. In fact, it provides the hydrodynamic equation for the process in (1.9) for which the invariant measures are not explicitly known and which can be seen as a non-trivial perturbation of the exclusion process.

Remark 1.4 (More general jump rates). The results of this article remain true for rather arbitrary jump rates. The particular choice of $c_{\pm 1}$ made in (1.3) allows us to keep the presentation and the proofs at a reasonable level of technicality. In Section 3.2 we state our theorems in more generality and we show where and how the proof presented in Section 2 has to be adapted.

Remark 1.5 (Beyond the macroscopic speed). In this paper we focus on the macroscopic speed of $\{x_t; t \geq 0\}$. It is natural to ask about fluctuations and large deviations for our model. It turns out that these two further questions are very delicate due to the lack of knowledge of the equilibrium measures of the process ξ_t . In other words, we need a finer control on the properties of the additive functional of the process in (2.2). We plan to address these questions in future works. Another direction which we are currently working on are conservative systems of ballistic particles. In such a setting the rescaling of the environment naturally matches the rescaling for the random walk.

The rest of the paper is organized as follows. Section 2 is devoted to the proof of Theorems 1.2 and 1.3. The proofs are based on the so-called *entropy method* and it will be divided in the following three main steps. *Step 1:* In Section 2.1 we show that the analysis of the random walk can be reduced to the study of the additive functional $\int_0^t (1 - 2\xi_s^n(0)) ds$, and we prove Theorem 2.1, a so-called *local replacement lemma*. This theorem allows to approximate the additive functional above in terms of the empirical measure in (1.10). *Steps 2 and 3:* Sections 2.2 and 2.3 deal with two necessary topological issues to show a weak convergence result of the type of Theorem 1.3, namely, tightness of

the sequence of considered processes, and the characterizations of the corresponding limiting points, respectively. In Section 3, we discuss possible variants or generalizations of the model presented in Section 1.1, for which the same techniques can be adapted. In particular in Section 3.1, we show how to derive the same type of results when considering the so-called speed-change exclusion as a dynamic environment. While Sections 3.2 and 3.3 focus on general jump rates for the random walk.

2. The entropy method

For $\rho \in [0, 1]$, let us denote by ν_ρ the product Bernoulli measure of density ρ , that is, ν_ρ is a probability measure in Ω such that

$$\nu_\rho(\eta(z_1) = 1, \dots, \eta(z_\ell) = 1) = \rho^\ell$$

for any set $\{z_1, \dots, z_\ell\} \subseteq \mathbb{Z}$. The measures ν_ρ are invariant with respect to the evolution of the process $\{\eta_t; t \geq 0\}$. The same is no longer true for the environment as seen by the walk, that is, the measures ν_ρ are *not left invariant* by the evolution of $\{\xi_t; t \geq 0\}$. In fact, if $\alpha > \beta$, the process x_t likes to be around the interface between regions with majority of particles and regions with majority of holes. This fact is the main difficulty in order to prove Theorems 1.2 and 1.3. As we will see, it turns out that the measure ν_ρ is close to be invariant in some sense.

2.1. A local replacement lemma

Let $\{\xi_t^n; t \geq 0\}$ be the rescaled environment as seen by the walk. For simplicity, we fix $T > 0$ and we consider the evolution of the process ξ_t^n up to time T .

Let us fix some notation. Let $\mathbb{D}([0, T]; \Omega)$ be the space of càdlàg trajectories from $[0, T]$ to Ω . Denote by \mathbb{P}^n the distribution in $\mathbb{D}([0, T]; \Omega)$ of the process $\{\xi_t^n; t \in [0, T]\}$ (with initial distribution μ^n) and by \mathbb{E}^n the expectation with respect to \mathbb{P}^n . We denote by \mathbb{P}_ρ^n the distribution of the process $\{\xi_t^n; t \in [0, T]\}$ starting from the distribution ν_ρ and by \mathbb{E}_ρ^n the expectation with respect to \mathbb{P}_ρ^n . We will use the same notations for the process $\{(\eta_t^n, x_t^n); t \in [0, T]\}$ with $x_0^n = 0$. Positive unspecified constants will be denoted by c .

We can recover the position x_t^n of the random walk in dynamical random environment looking at the signed number of shifts of the process $\{\xi_t^n; t \geq 0\}$ up to time t . Since the number of shifts is a compound Poisson process, we can write

$$\frac{x_t^n}{n} = \widetilde{M}_t^n + (\beta - \alpha) \int_0^t (1 - 2\xi_s^n(0)) \, ds, \tag{2.1}$$

where \widetilde{M}_t^n is a martingale of quadratic variation $\langle \widetilde{M}_t^n \rangle = \frac{(\alpha + \beta)t}{n}$. In particular, $\sup_{t \leq T} |\widetilde{M}_t^n|$ converges to 0 in probability as $n \rightarrow \infty$ for any $T \geq 0$. Therefore, in order to obtain a law of large numbers for the process $\{x_t^n; t \geq 0\}$ it is enough to obtain a law of large numbers for the integral

$$\int_0^t (1 - 2\xi_s^n(0)) \, ds. \tag{2.2}$$

What we will first prove is the following theorem.

Theorem 2.1 (Local replacement lemma). *Under the assumptions of Theorem 1.3,¹*

$$\lim_{\epsilon \rightarrow 0} \limsup_{n \rightarrow \infty} \mathbb{E}^n \left[\left| \int_0^t \left(\xi_s^n(0) - \frac{1}{\epsilon n} \sum_{z=1}^{\epsilon n} \xi_s^n(z) \right) \, ds \right| \right] = 0. \tag{2.3}$$

¹Here and below, for $\epsilon > 0$ and $n \in \mathbb{N}$, we use ϵn indistinctly for the real number ϵn and for its integer part $\lfloor \epsilon n \rfloor$.

This theorem is called *local replacement lemma* since allows to “replace” the local function $\xi_s^n(0)$ by the spatial average in (2.3) which can be rewritten in terms of the empirical measure $\hat{\pi}_s^n \in \mathcal{M}_+(\mathbb{R})$ in (1.10) as

$$\frac{1}{\epsilon n} \sum_{z=1}^{\epsilon n} \xi_s^n(z) = \int \epsilon^{-1} \mathbb{1}_{(0,\epsilon]}(x) \hat{\pi}_s^n(dx).$$

Such a replacement is a crucial step into the proof of the hydrodynamic limit of general diffusive particle systems and in fact, we will use it in the final argument of Section 2.3 to get our main theorems. Let us remark that a similar statement is proved in [10] in the context of a zero-range process as seen by a tagged particle. It turns out that the process considered in [10] has an invariant measure of product form, and by this reason the proof does not apply straightforwardly to our model. We will see below that an estimate on the Dirichlet form (see Lemma 2.2 below) is all that we need in order to adapt the proof of [10] to our setting and prove Theorem 2.1. The motivation for estimating the Dirichlet form is the following. When a Markov chain has a unique invariant measure, the (relative) entropy of the distribution of the chain with respect to the invariant measure is decreasing and vanishing in time. The speed at which the entropy decreases can be controlled by the Dirichlet form associated to the invariant measure. The same is true for general Markov processes, although the entropy with respect to some invariant measure may not go to zero, it is still decreasing, and the speed is still controlled by the associated Dirichlet form. For the process $\{\xi_t; t \geq 0\}$, the measure ν_ρ is not invariant, and therefore the entropy with respect to ν_ρ is not necessarily decreasing. This reflects on the fact that $\langle \sqrt{f}, \mathcal{L}_n \sqrt{f} \rangle$ may not be negative. However, in our model the entropy does not grow too much. In fact, under the assumptions of Theorem 1.2, as we show at the beginning of the proof of Theorem 2.1 below, the entropy between μ^n and ν_ρ is of order n . According to the entropy method introduced in [9], this entropy bound should be enough in order to prove the *replacement lemma*, and this is what we do next. We first introduce the Dirichlet form and prove the mentioned estimate in Lemma 2.2, then by using this estimate we prove Theorem 2.1.

Fix $\rho \in (0, 1)$, which we assume to be the one in the assumptions of Theorem 1.2. Let $f : \Omega \rightarrow \mathbb{R}$ be a density with respect to ν_ρ , that is, $f(\eta) \geq 0$ for every $\eta \in \Omega$ and $\int f d\nu_\rho = 1$. We define the *Dirichlet form* of f associated to \mathcal{L}^{ex} (see (1.9)) with respect to ν_ρ as

$$\mathcal{D}(f) = \frac{1}{2} \sum_{z \in \mathbb{Z}} \int (\sqrt{f(\eta^{z,z+1})} - \sqrt{f(\eta)})^2 \nu_\rho(d\eta). \quad (2.4)$$

Let us denote by $\langle \cdot, \cdot \rangle$ the inner product in $L^2(\nu_\rho)$. It is standard to check that

$$\mathcal{D}(f) = \langle \sqrt{f}, -\mathcal{L}^{\text{ex}} \sqrt{f} \rangle. \quad (2.5)$$

We now show a crucial bound on $\langle \sqrt{f}, \mathcal{L}_n \sqrt{f} \rangle$.

Lemma 2.2 (Dirichlet form estimate). *For any density $f : \Omega \rightarrow \mathbb{R}$ with respect to ν_ρ ,*

$$\langle \sqrt{f}, \mathcal{L}_n \sqrt{f} \rangle \leq -n^2 \mathcal{D}(f) + 2n|\alpha - \beta|.$$

Proof. As in equation (1.9), we can split \mathcal{L}_n into two parts: $n^2 \mathcal{L}^{\text{ex}}$, which is the rescaled part of the generator corresponding to jumps of the particles, and $n \mathcal{L}^{\text{rw}}$, which corresponds to rescaled jumps of the random walk. By (2.5), the part of the generator corresponding to jumps of the particles satisfies $\langle \sqrt{f}, n^2 \mathcal{L}^{\text{ex}} \sqrt{f} \rangle = -n^2 \mathcal{D}(f)$. Therefore, we just need to prove that

$$\langle \sqrt{f}, \mathcal{L}^{\text{rw}} \sqrt{f} \rangle \leq 2|\alpha - \beta|$$

for any density f . Notice that $\langle \sqrt{f}, \sqrt{f} \rangle = 1$ and notice that also $\langle \sqrt{\theta^1 f}, \sqrt{\theta^1 f} \rangle = 1$. This last identity follows from the invariance of the measure ν_ρ under spatial shifts. By the same reason, $\langle \sqrt{\theta^{-1} f}, \sqrt{\theta^{-1} f} \rangle = 1$. By the Cauchy–Schwarz inequality,

$$\langle \sqrt{f}, \sqrt{\theta^1 f} \rangle \leq \frac{1}{2} (\langle \sqrt{f}, \sqrt{f} \rangle + \langle \sqrt{\theta^1 f}, \sqrt{\theta^1 f} \rangle) = 1.$$

Therefore,

$$\begin{aligned} & \langle \sqrt{f}, (\beta + (\alpha - \beta)\xi(0))(\sqrt{\theta^1 f} - \sqrt{f}) \rangle \\ & \leq \max\{\beta + (\alpha - \beta)\xi(0)\} \langle \sqrt{f}, \sqrt{\theta^1 f} \rangle \\ & \quad - \min\{\beta + (\alpha - \beta)\xi(0)\} \langle \sqrt{f}, \sqrt{f} \rangle \\ & \leq \max\{\beta + (\alpha - \beta)\xi(0)\} - \min\{\beta + (\alpha - \beta)\xi(0)\} \\ & \leq |\alpha - \beta|. \end{aligned}$$

The same reasoning allows to bound the other term in $\langle \sqrt{f}, \mathcal{L}^{\text{rw}} \sqrt{f} \rangle$, see (1.9), which ends the proof of the lemma. \square

We are now in shape to prove Theorem 2.1, using Lemma 2.2.

Proof of Theorem 2.1. Let us denote by $H(\mu|\nu)$ the entropy between two given measures. First we observe that $H(\mathbb{P}^n|\mathbb{P}_\rho^n) = H(\mu^n|\nu_\rho) \leq cn$. The first equality is an easy general fact about Markov chains. The second inequality follows from the structure of the measures μ^n and ν_ρ and the assumption in (1.7). By the entropy inequality, see Proposition 8.1 in [11], for any function $V : \Omega \rightarrow \mathbb{R}$ and any $\gamma > 0$,

$$\mathbb{E}^n \left[\left| \int_0^t V(\xi_s^n) ds \right| \right] \leq \frac{H(\mathbb{P}^n|\mathbb{P}_\rho^n)}{\gamma n} + \frac{1}{\gamma n} \log \mathbb{E}_\rho^n [e^{\gamma n |\int_0^t V(\xi_s^n) ds|}]. \tag{2.6}$$

The first term on the right-hand side of (2.6) is then bounded by $\frac{c}{\gamma}$. Using the estimate $e^{|a|} \leq e^a + e^{-a}$, we can get rid of the absolute value on the second term of the right-hand side of (2.6), at the cost of estimating two expectations, one involving V and another involving $-V$. Therefore, it suffices to show that

$$\lim_{\gamma \rightarrow \infty} \limsup_{\epsilon \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{1}{\gamma n} \log \mathbb{E}_\rho^n [e^{\gamma n \int_0^t V(\xi_s^n) ds}] = 0, \tag{2.7}$$

where

$$V(\xi) = \pm \left(\xi(0) - \frac{1}{\epsilon n} \sum_{z=1}^{\epsilon n} \xi(z) \right).$$

By Feynman–Kac formula, we can express the expectation in (2.7) in terms of the semigroup associated to the operator $\mathcal{L}_n + \gamma n V$. After some computations based on spectral theory (see Lemma A.1.7.2 in [11]), we can obtain the bound

$$\frac{1}{\gamma n} \log \mathbb{E}_\rho^n [e^{\gamma n \int_0^t V(\xi_s^n) ds}] \leq t \sup_f \left\{ \langle V, f \rangle + \frac{1}{\gamma n} \langle \sqrt{f}, \mathcal{L}_n \sqrt{f} \rangle \right\},$$

where the supremum is taken over all the densities f with respect to ν_ρ . By Lemma 2.2, the supremum above is bounded by

$$\frac{2t|\alpha - \beta|}{\gamma} + t \sup_f \left\{ \langle V, f \rangle - \frac{n}{\gamma} \mathcal{D}(f) \right\}.$$

Therefore, in order to prove the theorem, we only need to prove that

$$\lim_{\gamma \rightarrow \infty} \limsup_{\epsilon \rightarrow 0} \limsup_{n \rightarrow \infty} \sup_f \left\{ \langle V, f \rangle - \frac{n}{\gamma} \mathcal{D}(f) \right\} = 0. \tag{2.8}$$

Notice that

$$\xi(0) - \frac{1}{\epsilon n} \sum_{z=1}^{\epsilon n} \xi(z) = \sum_{z=1}^{\epsilon n} \frac{\epsilon n - z + 1}{\epsilon n} (\xi(z-1) - \xi(z)).$$

We first estimate $\langle \xi(z-1) - \xi(z), f \rangle$. Performing the change of variables $\xi \rightarrow \xi^{z,z+1}$, we see that

$$\langle \xi(z-1) - \xi(z), f \rangle = \int \xi(z) (f(\xi^{z,z+1}) - f(\xi)) \nu_\rho(d\xi).$$

Let us define

$$\mathcal{D}_z(f) = \int (\sqrt{f(\xi^{z,z+1})} - \sqrt{f(\xi)})^2 \nu_\rho(d\xi),$$

in such a way that $\mathcal{D}(f) = \frac{1}{2} \sum_z \mathcal{D}_z(f)$. Writing

$$f(\xi^{z,z+1}) - f(\xi) = \{\sqrt{f(\xi^{z,z+1})} - \sqrt{f(\xi)}\} \{\sqrt{f(\xi^{z,z+1})} + \sqrt{f(\xi)}\},$$

using the Young inequality and the fact that $\int f d\nu_\rho = 1$ and $0 \leq \xi(z) \leq 1$, we see that

$$\langle \xi(z-1) - \xi(z), f \rangle \leq \frac{1}{2\lambda_z} \mathcal{D}_z(f) + 2\lambda_z$$

for any $\lambda_z > 0$. Choosing

$$\lambda_z = \frac{\gamma}{n} \frac{\epsilon n - z + 1}{\epsilon n},$$

we obtain

$$\langle V, f \rangle - \frac{n}{\gamma} \mathcal{D}(f) \leq \frac{2\gamma}{\epsilon^2 n^3} \sum_{z=1}^{\epsilon n} (\epsilon n - z + 1)^2 \leq c\gamma\epsilon$$

for any density f , which gives (2.8) and concludes the theorem. □

2.2. Tightness

The proof of a theorem like Theorem 1.3 is usually performed following the classical three-steps procedure to prove weak convergence of probability measures in Polish spaces. The first step is to prove tightness of the corresponding sequence of distributions with respect to some properly chosen topology. The second step is to prove that any limit point (they exists due to tightness and Prohorov’s Theorem) satisfies a convenient set of properties. The third step is to prove that there exists at most one probability measure on the corresponding Polish space satisfying those properties. Then, an abstract topology result, namely that any relatively compact set with a unique limit point is a converging sequence shows that the sequence of distributions converges in distribution with respect to the already chosen topology. In the case of the hydrodynamic limit stated in Theorem 1.3, convergence in probability is readily obtained, since the limit is non-random. In this section we show the first of these steps: the tightness. Before entering into the details, we do a small detour to explain the choice of topology and some facts associated to this choice.

Let $\mathcal{B}(\mathbb{R})$ denote the family of Borel sets of \mathbb{R} , that is, the smallest σ -algebra formed by subsets of \mathbb{R} containing all the open sets of \mathbb{R} . We say that a measure $\pi(dx)$ defined in \mathbb{R} is *non-negative* if $\pi(A) \geq 0$ for any $A \in \mathcal{B}(\mathbb{R})$. We say that the measure $\pi(dx)$ is a *Radon* measure if $-\infty < \pi(K) < +\infty$ for any compact set $K \subseteq \mathbb{R}$. The empirical measure $\{\hat{\pi}_t^n(dx); t \in [0, T]\}$, is a random, non-negative, Radon measure. Let us denote by $\mathcal{M}_+(\mathbb{R})$ the set of non-negative, Radon measures in \mathbb{R} . The weak topology in $\mathcal{M}_+(\mathbb{R})$ is defined in the following way. Let $\mathcal{C}_c(\mathbb{R})$ denote the set of

functions $f : \mathbb{R} \rightarrow \mathbb{R}$ which are continuous and of bounded support. We say that a sequence $\{\pi^n; n \in \mathbb{N}\}$ in $\mathcal{M}_+(\mathbb{R})$ converges to a measure $\pi \in \mathcal{M}_+(\mathbb{R})$ if for any $f \in \mathcal{C}_c(\mathbb{R})$,

$$\lim_{n \rightarrow \infty} \int f \, d\pi^n = \int f \, d\pi.$$

The space $\mathcal{M}_+(\mathbb{R})$ turns out to be a Polish space with respect to the weak topology. In fact, there exists a sequence $\{f_\ell; \ell \in \mathbb{N}\}$ of functions in $\mathcal{C}_c(\mathbb{R})$ such that the distance $d : \mathcal{M}_+(\mathbb{R}) \times \mathcal{M}_+(\mathbb{R})$ defined by

$$d(\pi, \pi') = \sum_{\ell \in \mathbb{N}} \frac{1}{2^\ell} \min \left\{ \left| \int f_\ell \, d(\pi - \pi') \right|, 1 \right\}$$

is a metric for the weak topology in $\mathcal{M}_+(\mathbb{R})$. We can assume that the functions $\{f_\ell; \ell \in \mathbb{N}\}$ are infinitely differentiable. We refer the reader to Section 4 of [11] for the latter statement as well as for the following topological considerations. In order to simplify the notation, we will write $\pi(f) = \int f \, d\pi$ for $\pi \in \mathcal{M}_+(\mathbb{R})$ and $f \in \mathcal{C}_c(\mathbb{R})$. Let us consider the following topology in $\mathcal{C}_c(\mathbb{R})$. We say that the sequence $\{f_n; n \in \mathbb{N}\}$ of functions in $\mathcal{C}_c(\mathbb{R})$ converges to $f \in \mathcal{C}_c(\mathbb{R})$ if two things happen:

- (i) there exists a compact K such that the support of f_n is contained in K for any $n \in \mathbb{N}$,
- (ii) $\lim_{n \rightarrow \infty} \sup_{x \in \mathbb{R}} |f_n(x) - f(x)| = 0$.

It turns out that $\mathcal{M}_+(\mathbb{R})$ is the dual of $\mathcal{C}_c(\mathbb{R})$ with respect to this topology, and moreover, the weak topology in $\mathcal{M}_+(\mathbb{R})$ is the weak- $*$ topology of $\mathcal{M}_+(\mathbb{R})$ associated to this duality. This fact provides us with a very simple tightness criterion for measure-valued processes:

Proposition 2.3. *Let $\{\pi_t^n(dx); t \in [0, T]\}_{n \in \mathbb{N}}$ be a sequence of measure-valued processes with trajectories in the space $\mathbb{D}([0, T]; \mathcal{M}_+(\mathbb{R}))$. The sequence of processes $\{\pi_t^n(dx); t \in [0, T]\}_{n \in \mathbb{N}}$ is tight with respect to the J_1 -Skorohod topology of $\mathbb{D}([0, T]; \mathcal{M}_+(\mathbb{R}))$ if and only if the sequence of real-valued processes $\{\pi_t^n(f_\ell); t \in [0, T]\}_{n \in \mathbb{N}}$ is tight with respect to the J_1 -Skorohod topology of $\mathbb{D}([0, T]; \mathbb{R})$ for any $\ell \in \mathbb{N}$. If, for any $\ell \in \mathbb{N}$, any limit point of $\{\pi_t^n(f_\ell); t \in [0, T]\}_{n \in \mathbb{N}}$ is supported on the space $\mathcal{C}([0, T]; \mathbb{R})$ of continuous functions, then any limit point of $\{\pi_t^n(dx); t \in [0, T]\}_{n \in \mathbb{N}}$ is supported on the space $\mathcal{C}([0, T]; \mathcal{M}_+(\mathbb{R}))$.*

Roughly speaking, this proposition is saying that the verification of tightness for measure-valued processes can be reduced to the verification of tightness for real-valued processes. This property holds true for any dual of a Polish space equipped with the weak- $*$ topology. For a proof of this proposition in the context of measure-valued processes, see Section 4 of [11]. The proof can be easily adapted to any dual of a Polish space.

Now we are in position to state the tightness results for the random walk and the environment as viewed by the walk:

Theorem 2.4. *The sequence of processes $\{\hat{\pi}_t^n; t \in [0, T]\}_{n \in \mathbb{N}}$ is tight with respect to the J_1 -Skorohod topology of $\mathbb{D}([0, T]; \mathcal{M}_+(\mathbb{R}))$. Moreover, any limit point of $\{\hat{\pi}_t^n; t \in [0, T]\}_{n \in \mathbb{N}}$ is supported on the set $\mathcal{C}([0, T]; \mathcal{M}_+(\mathbb{R}))$ of continuous trajectories. The sequence of real-valued processes $\{n^{-1}x_t^n; t \in [0, T]\}_{n \in \mathbb{N}}$ is tight with respect to the J_1 -Skorohod topology of $\mathbb{D}([0, T]; \mathbb{R})$. Moreover, any limit point of $\{n^{-1}x_t^n; t \in [0, T]\}_{n \in \mathbb{N}}$ is supported on the set $\mathcal{C}([0, T]; \mathbb{R})$.*

Proof. Let us recall Dynkin’s formula for a function of a Markov process: for any local function $F : \Omega \rightarrow \mathbb{R}$, the process

$$F(\xi_t^n) - F(\xi_0^n) - \int_0^t \mathcal{L}_n F(\xi_s^n) \, ds$$

is a mean-zero martingale. The quadratic variation of this martingale can also be computed in terms of the generator \mathcal{L}_n and it is given by

$$\int_0^t \mathcal{L}_n F(\xi_s^n)^2 - 2F(\xi_s^n)\mathcal{L}_n F(\xi_s^n) \, ds = \int_0^t [\mathcal{L}_n(F - F(\xi_s^n))^2](\xi_s^n) \, ds.$$

Let $f \in \mathcal{C}_c(\mathbb{R})$ be a smooth function. Let us define the discrete gradient(s) and discrete Laplacian of f by

$$\begin{aligned}\Delta_n f\left(\frac{x}{n}\right) &= n^2 \left[f\left(\frac{x+1}{n}\right) + f\left(\frac{x-1}{n}\right) - 2f\left(\frac{x}{n}\right) \right], \\ \nabla_-^n f\left(\frac{x}{n}\right) &= n \left[f\left(\frac{x}{n}\right) - f\left(\frac{x-1}{n}\right) \right], \\ \nabla_+^n f\left(\frac{x}{n}\right) &= n \left[f\left(\frac{x+1}{n}\right) - f\left(\frac{x}{n}\right) \right].\end{aligned}$$

Taking $F(\xi_t^n) = \hat{\pi}_t^n(f)$, we see that

$$\begin{aligned}M_t^n(f) &= \hat{\pi}_t^n(f) - \hat{\pi}_0^n(f) - \int_0^t \left\{ \hat{\pi}_s^n(\Delta_n f) - (\beta - \alpha)(1 - 2\xi_s^n(0))\hat{\pi}_s^n(\nabla_-^n f) \right. \\ &\quad \left. - \frac{1}{n}[\alpha + (\beta - \alpha)\xi_s^n(0)]\hat{\pi}_s^n(\Delta_n f) \right\} ds\end{aligned}$$

is a martingale. Note that the last term comes from the difference between $\nabla_+^n f - \nabla_-^n f$. Its quadratic variation is

$$\begin{aligned}\langle M_t^n(f) \rangle &= \frac{1}{n} \int_0^t \frac{1}{n} \sum_{x \in \mathbb{Z}} (\xi_s^n(x) - \xi_s^n(x-1))^2 \left(\nabla_-^n f\left(\frac{x}{n}\right) \right)^2 \\ &\quad + \frac{\alpha + \beta}{n} \int_0^t (\hat{\pi}_s^n(\nabla_-^n f))^2 ds + E_n,\end{aligned}$$

where E_n is a lower order term which captures the influence of $\nabla_+^n f - \nabla_-^n f$ and is given by

$$\frac{1}{n} \int_0^t (\beta \xi_s^n(0) + \alpha(1 - \xi_s^n(0))) [(\hat{\pi}_s^n(\nabla_+^n f))^2 - (\hat{\pi}_s^n(\nabla_-^n f))^2] ds.$$

Notice that the occupation variables can only assume the values 0, 1. Since f is infinitely differentiable and of compact support, there exists a constant $C(f) > 0$ such that

$$|\langle M_t^n(f) \rangle| \leq \frac{C(f)t}{n}$$

for any $t \in [0, T]$ and any $n \in \mathbb{N}$. Note that $\mathbb{E}^n[M_t^n(f)^2] = \mathbb{E}^n[\langle M_t^n(f) \rangle]$. Therefore, by Doob's inequality we have

$$\mathbb{P}^n \left[\sup_{0 \leq t \leq T} |M_t^n(f)| \geq \epsilon \right] \leq \frac{4C(f)T}{\epsilon^2 n}$$

for any $n \in \mathbb{N}$. We conclude that the process $\{M_t^n(f); t \geq 0\}$ converges to 0 in probability (and therefore in distribution) with respect to the uniform topology in $\mathbb{D}([0, T]; \mathbb{R})$. Notice that $\hat{\pi}_0^n(f) = \pi_0^n(f)$. Therefore, by hypothesis $\hat{\pi}_0^n(f)$ converges in probability (and in distribution) to $\int u_0(x) f(x) dx$. These two convergences reduce the proof of tightness of $\{\hat{\pi}_t^n(f); t \in [0, T]\}$ to the proof of tightness of the integral term

$$\mathcal{I}_t^n(f) = \int_0^t \left\{ \hat{\pi}_s^n(\Delta_n f) - (\beta - \alpha)(1 - 2\xi_s^n(0))\hat{\pi}_s^n(\nabla_-^n f) \right\} ds.$$

But, using again the boundedness of $\xi_s^n(0)$,

$$|\mathcal{I}_t^n(f) - \mathcal{I}_s^n(f)| \leq \frac{|t-s|}{n} \sum_{x \in \mathbb{Z}} \left(\left| \Delta_n f\left(\frac{x}{n}\right) \right| + \left| \nabla_-^n f\left(\frac{x}{n}\right) \right| \right),$$

and by the smoothness of f we conclude that the sequence of processes $\{\mathcal{I}_t^n; t \in [0, T]\}_{n \in \mathbb{N}}$ is uniformly Lipschitz (uniformly in n and t), and in particular it is tight with respect to the uniform topology of $\mathcal{C}([0, T]; \mathbb{R})$. This proves two things: first, the sequence $\{\hat{\pi}_t^n(f); t \in [0, T]\}_{n \in \mathbb{N}}$ is tight; and second, every limit point of $\{\hat{\pi}_t^n(f); t \in [0, T]\}_{n \in \mathbb{N}}$ is supported on continuous trajectories. By Proposition 2.3, the sequence of measure-valued processes $\{\hat{\pi}_t^n; t \in [0, T]\}_{n \in \mathbb{N}}$ is tight with respect to the J_1 -Skorohod topology on $\mathbb{D}([0, T]; \mathcal{M}_+(\mathbb{R}))$ and any limit point is supported on continuous trajectories.

Now we turn into the proof of tightness of the sequence $\{\frac{x_t^n}{n}; t \in [0, T]\}$. This is actually simpler. In fact, the process

$$\widetilde{M}_t^n := \frac{x_t^n}{n} - (\beta - \alpha) \int_0^t (1 - 2\xi_s^n(0)) ds$$

is a martingale of quadratic variation $\frac{(\alpha+\beta)t}{n}$. As above, by Doob’s inequality the martingale converges to 0 in probability with respect to the uniform topology on $\mathbb{D}([0, T]; \mathbb{R})$. Again as above, the integral term is uniformly Lipschitz, both in t and n . In fact, the Lipschitz constant is bounded above by $|\beta - \alpha|$. Therefore, $\{\frac{x_t^n}{n}; t \in [0, T]\}$ is tight with respect to the J_1 -Skorohod topology on $\mathbb{D}([0, T]; \mathbb{R})$ and any limit point is supported on continuous trajectories. In fact, we can say a little bit more about the limit points: they are supported on *Lipschitz functions* of Lipschitz constant bounded by $|\beta - \alpha|$. □

2.3. Characterization of limit points: Proofs of Theorems 1.2 and 1.3

We are now ready to finish our proofs. We first observe that a vector of random processes is tight if and only if each coordinate is tight. Therefore, by Theorem 2.4, the triple $\{(\pi_t^n, \hat{\pi}_t^n, \frac{x_t^n}{n}); t \in [0, T]\}_{n \in \mathbb{N}}$ is tight. Let $\{(u(t, x) dx, \hat{u}_t(dx), f(t)); t \in [0, T]\}$ be a limit point of the triple and let n' be the subsequence of n for which the triple converges to that limit point. Remind that we already know that $u(t, x) dx$ is the solution of the heat equation, and in particular it is deterministic, but in principle $f(t)$ and $\hat{u}_t(dx)$ may be random. A first observation is that $\hat{u}_t(dx)$ has a density with respect to Lebesgue measure. This is an easy consequence of the boundedness of ξ_s^n . In fact, for any closed, non-empty interval $A \subseteq \mathbb{R}$, $\hat{\pi}_t^n(\mathbb{1}_A) \leq |A| + \frac{1}{n}$, where $|A|$ denotes the Lebesgue measure of A and $\mathbb{1}$ is the usual characteristic function. Therefore, $\hat{u}_t(dx) = \hat{u}(t, x) dx$ for some random function $\{\hat{u}(t, x); t \in [0, T], x \in \mathbb{R}\}$, bounded between 0 and 1. Another observation is that $\{f(t); t \in [0, T]\}$ is Lipschitz, with Lipschitz constant bounded above by $|\beta - \alpha|$. For any fixed $n \in \mathbb{N}$, the measure $\hat{\pi}_t^n$ is the shift of π_t^n by $\frac{x_t^n}{n}$. Since shifting by a continuous function is a continuous operation on the space $\mathcal{M}_+(\mathbb{R})$, this relation is also satisfied by the limiting processes. Therefore, we have the relation $\hat{u}(t, x) = u(t, x + f(t))$ for any $t \in [0, T]$ and any $x \in \mathbb{R}$.

Next, recall that the sum involved in equation (2.3) is equal to $\hat{\pi}_s^n(\epsilon^{-1} \mathbb{1}_{(0, \epsilon]})$. Since the step function $\epsilon^{-1} \mathbb{1}_{(0, \epsilon]}$ is not a continuous function, we cannot say that $\hat{\pi}_s^n(\epsilon^{-1} \mathbb{1}_{(0, \epsilon]})$ converges. However, since the limiting measure $\hat{u}(s, x) dx$ has a density and the step function $\epsilon^{-1} \mathbb{1}_{(0, \epsilon]}$ is *a.s.* continuous, we have that

$$\lim_{n \rightarrow \infty} \hat{\pi}_s^n(\epsilon^{-1} \mathbb{1}_{(0, \epsilon]}) = \epsilon^{-1} \int_0^\epsilon \hat{u}(s, x) dx.$$

On the other hand, by (2.1),

$$\int_0^t (1 - 2\xi_s^n(0)) ds = \frac{1}{\beta - \alpha} \left(\frac{x_t^n}{n} - \widetilde{M}_t^n \right),$$

which converges to $(\beta - \alpha)^{-1} f(t)$. Replacing these two limits into (2.3), and using the identity $\hat{u}(t, x) = u(t, x + f(t))$ we obtain the relation

$$\lim_{\epsilon \rightarrow 0} \mathbb{E} \left[\left| f(t) - \frac{\beta - \alpha}{\epsilon} \int_0^t \int_0^\epsilon (1 - 2u(s, x + f(s))) dx ds \right| \right] = 0.$$

Since f is uniformly Lipschitz and u is smooth, the limit as $\epsilon \rightarrow 0$ of the integral above is equal to

$$(\beta - \alpha) \int_0^t (1 - 2u(s, f(s))) ds.$$

We conclude that $\{f(t); t \in [0, T]\}$ satisfies the integral equation

$$f(t) = (\beta - \alpha) \int_0^t (1 - 2u(s, f(s))) ds,$$

which is nothing but the integral version of (1.8). Since this equation has a unique solution, we conclude that f is deterministic and uniquely defined. This ends the proof of Theorem 1.2. Theorem 1.3 follows from the argument, after recalling that $\hat{u}(t, x) = u(t, x + f(t))$ for all $x \in \mathbb{R}$, in particular $x = 0$.

3. Generalizations

In this section, we present some generalizations of Theorems 1.2 and 1.3. One possibility is to extend the results for other underlying dynamics. This is the content of Section 3.1. A second one is to consider different transitions for the random walk. In Sections 3.2 and 3.3 we discuss how to generalize Theorems 1.2 and 1.3 for random walks with more general jump rates and with macroscopic jumps, respectively.

3.1. The speed-change exclusion process

An example of underlying dynamics for which we can extend our results in a straightforward way is the so-called *speed-change* exclusion process, or stochastic lattice gas at infinite temperature. In this dynamics, exchanges of particles between sites z and $z + 1$ are performed at rate $c_z(\eta) = c_0(\theta^{-z}\eta)$, where $c_0(\eta)$ is a strictly positive, local function which does not depend on the values of $\eta(0)$ and $\eta(1)$. More precisely, let $c_0 : \Omega \rightarrow \mathbb{R}$ be a local, positive function. Notice that positivity plus locality imply that there exists a constant $\epsilon_0 > 0$ such that $\epsilon_0 \leq c_0(\eta) \leq \epsilon_0^{-1}$ for any $\eta \in \Omega$. The generator of the speed-change simple exclusion process acts on local functions $f : \Omega \rightarrow \mathbb{R}$ as

$$\mathcal{L}^{\text{sc}} f(\eta) = \sum_{z \in \mathbb{Z}} c_z(\eta) \{f(\eta^{z,z+1}) - f(\eta)\}, \tag{3.1}$$

where $c_z(\eta) = c_0(\theta^{-z}\eta)$. The stochastic evolution can be described as follows. At rate $c_z(\eta)$ the occupation variables $\eta(z), \eta(z + 1)$ are exchanged, and this rate depends on the occupation of the neighbors of $z, z + 1$ up to some finite distance R . Such dependency gives raise to a non-linearity in the hydrodynamic limit. For this model, under the assumptions of Proposition 1.1, the statement holds true with a Cauchy problem of the form

$$\begin{cases} \partial_t u(t, x) = (1/2) \partial_{xx}^2 \Phi(u(t, x)), & t \geq 0, x \in \mathbb{R}, \\ u(0, x) = u_0(x), & x \in \mathbb{R}, \end{cases}$$

where $\Phi(\rho)$ is given in general by a variational formula (see Section 7 of [11] for more details).

A choice which is very popular in the literature is $c_x(\eta) = 1 + a(\eta(x - 1) + \eta(x + 2))$ for some $a > -\frac{1}{2}$. For this particular choice, the model turns out to be gradient, see, e.g., [8], and $\Phi(\rho) = \rho + a\rho^2$.

The generator of the environment as seen by the walker x_t can be written as $\tilde{\mathcal{L}}_n = n^2 \mathcal{L}^{\text{sc}} + n \mathcal{L}^{\text{rw}}$, with \mathcal{L}^{sc} as in equation (3.1). We have the following version of Lemma 2.2:

Lemma 3.1. *For any density $f : \Omega \rightarrow \mathbb{R}$,*

$$\langle \sqrt{f}, \mathcal{L}_n \sqrt{f} \rangle \leq -\frac{n^2}{\epsilon_0} \mathcal{D}(f) + 2n|\alpha - \beta|.$$

Keeping this lemma in mind, the proof of the replacement lemma, Theorem 2.1, can be repeated for the speed-change exclusion process. *Mutatis mutandis*, the rest of the proofs is the same, as well the conclusions in the Theorem 1.2 and the Theorem 1.3.

3.2. Random walks with more general jump rates

For simplicity, the random walk we defined in Section 1.1 looks only at the state of the exclusion process at its current position. However, our result holds for a more general choice of the transition rates. Let $\gamma_z : \Omega \rightarrow [0, \infty)$, $z \in \mathbb{Z}$, be a collection of local functions prescribing the jump rates of the random walk, i.e.,

$$Lf(\eta, x) = \sum_{z \in \mathbb{Z}} [f(\eta^{z, z+1}, x) - f(\eta, x)] + \sum_{z \in \mathbb{Z}} \gamma_z(\theta^{-x} \eta) [f(\eta, x+z) - f(\eta, x)]. \tag{3.2}$$

Note that the setting discussed in the bulk of the paper corresponds to $\gamma_{+1}(\eta) = \beta + (\alpha - \beta)\eta(0)$, $\gamma_{-1}(\eta) = \alpha + (\beta - \alpha)\eta(0)$, and $\gamma_z(\eta) = 0$ else.

In the general case described in (3.2), by assuming that $\sum_{z \in \mathbb{Z}} \sup_{\eta} \tilde{\gamma}_z(\eta) < \infty$, the hydrodynamic limit of Theorem 1.3 still holds with $\hat{u}(t, x)$ being the solution of

$$\begin{cases} \partial_t \hat{u}(t, x) = (1/2) \partial_{xx}^2 \hat{u}(t, x) + \gamma(\hat{u}(t, 0)) \partial_x \hat{u}(t, x), & t \geq 0, x \in \mathbb{R}, \\ \hat{u}(0, x) = u_0(x), & x \in \mathbb{R}, \end{cases}$$

where $\gamma(\rho) = \nu_{\rho}(\sum_z z \gamma_z)$.

In fact, with $V = (\xi(0) - \frac{1}{\epsilon n} \sum_{x=1}^{\epsilon n} \xi(x))$ replaced by $V_{\epsilon n} = \sum_z z \gamma_z(\xi) - \gamma(\frac{1}{\epsilon n} \sum_{x=1}^{\epsilon n} \xi(x))$ the proofs generalize to this setting except for equation (2.8). Here a more sophisticated argument is necessary. The proof follows from the arguments in [10]. Since the reference [10] is quite technical, we try to be more specific. The proof of (2.8) follows the celebrated one-block, two-blocks scheme introduced by [9]. The one-block estimate reduces to prove that

$$\lim_{\gamma \rightarrow \infty} \limsup_{\ell \rightarrow \infty} \limsup_{n \rightarrow \infty} \sup_f \left\{ \langle V_{\ell}, f \rangle - \frac{n}{\gamma} \mathcal{D}(f) \right\} = 0.$$

This is equivalent to the estimation of equation (6.2) in [10]. The two-blocks estimate reduces to prove that

$$\lim_{\gamma \rightarrow \infty} \limsup_{\ell \rightarrow \infty} \limsup_{\epsilon \rightarrow 0} \limsup_{n \rightarrow \infty} \sup_{f, x} \left\{ \langle V_{\ell, x}, f \rangle - \frac{n}{\gamma} \mathcal{D}(f) \right\} = 0,$$

where

$$V_{\ell, x}(\xi) = \gamma \left(\frac{1}{\ell} \sum_{y=1}^{\ell} \xi(y) \right) - \gamma \left(\frac{1}{\ell} \sum_{y=x+1}^{x+\ell} \xi(y) \right)$$

and the supremum is over densities $f : \Omega \rightarrow \mathbb{R}$ and over $2\ell + 1 \leq x \leq \epsilon n$. This is basically what is proven in Lemma 6.5 of [10]. The rest of the proof follows like in the proof of Proposition 6.1 of [10].

Remark 3.2. One of the main assumptions in [10] is a sharp lower bound on the spectral gap of the dynamics restricted to a finite box, which is well-known for the exclusion process.

3.3. Random walks with macroscopic jumps

For the hydrodynamic limit as presented to hold it is necessary that the rates of the random walk are properly rescaled. It is however possible to introduce rare large-scale jumps. Rescaling (3.2), we get

$$L_n f(\eta, x) = n^2 \sum_{z \in \mathbb{Z}} [f(\eta^{z, z+1}, x) - f(\eta, x)] + n \sum_{z \in \mathbb{Z}} \gamma_z(\theta^{-x} \eta) [f(\eta, x+z) - f(\eta, x)].$$

We can introduce long range jumps by adding a third term:

$$L_n f(\eta, x) = n^2 \sum_{z \in \mathbb{Z}} [f(\eta^{z, z+1}, x) - f(\eta, x)] + n \sum_{z \in \mathbb{Z}} \gamma_z(\theta^{-x} \eta) [f(\eta, x+z) - f(\eta, x)]$$

$$+ \sum_{z \in \mathbb{Z}} \tilde{\gamma}_z(\theta^{-x} \eta) [f(\eta, x + nz) - f(\eta, x)], \quad (3.3)$$

with

$$\sum_{z \in \mathbb{Z}} \sup_{\eta} \tilde{\gamma}_z(\eta) < \infty. \quad (3.4)$$

Note how the jump rates $\tilde{\gamma}_z: \Omega \rightarrow [0, \infty)$, $z \in \mathbb{Z}$, are not rescaled in time, but the corresponding jumps are of order n . This leads to randomness in the hydrodynamic limit, where the random walk x_t^n converges to a space–time inhomogeneous random walk x_t on \mathbb{R} with drift, characterized by the generator

$$\begin{aligned} L_t^{\text{rw}} f(x) &= \sum_{z \in \mathbb{Z}} \tilde{\gamma}_z(u(t, x)) [f(x + z) - f(x)] + \gamma(u(t, x)) f'(x), \\ \tilde{\gamma}_z(\rho) &= \nu_\rho(\tilde{\gamma}_z). \end{aligned} \quad (3.5)$$

The idea of the proof is rather straightforward, using the hydrodynamic limit without long-range jumps. We now give the main lines of this proof.

Let τ^n be the time of the first macroscopic jump and z^n the jump size over n , i.e.,

$$x_{\tau^n}^n = x_{\tau^n-}^n + nz^n.$$

The time τ^n is distributed according to the first arrival of any of the Poisson point processes $\{N_z^n: z \in \mathbb{Z}\}$ on $[0, \infty)$ with intensity measure $\{I_z^n: z \in \mathbb{Z}\}$ given by

$$I_z^n([a, b]) = \int_a^b \tilde{\gamma}_z(\xi_t^n) dt.$$

From the hydrodynamic limit without the macroscopic jumps, we know that N_z^n converges to the limiting Poisson point process N_z with intensity measures

$$I_z([a, b]) = \int_a^b \tilde{\gamma}_z(\hat{u}(t, 0)) dt.$$

In Lemma 3.3 below, we prove that for $t \in [0, \tau^n]$, x_t^n converges to x_t , $t \in [0, \tau]$, where x_t is the random walk described by the generator L_t^{rw} in (3.5), and τ is the first macroscopic jump time, corresponding to the first arrival of the Poisson point processes $\{N_z: z \in \mathbb{Z}\}$.

Finally, the result follows by iterating the same argument for the other macroscopic jumps after the first.

Lemma 3.3. *Define $X_{[0, \tau^n]}^n := \{x_t^n: t \in [0, \tau^n]\}$ and $X_{[0, \tau]} := \{x_t: t \in [0, \tau]\}$. Then,*

$$X_{[0, \tau^n]}^n \text{ converges in distribution to } X_{[0, \tau]},$$

as n goes to infinity.

Proof. The idea is to couple the Poisson point processes $\{N_z: z \in \mathbb{Z}\}$ and $\{N_z^n: z \in \mathbb{Z}\}$. For $z \in \mathbb{Z}$, let $\bar{N}_z := (\bar{N}_z(t))_{t \geq 0}$ a Poisson process with rate $\|\tilde{\gamma}_z\|_\infty$. For $t \geq 0$, consider the time-changed process

$$M_z^n(t) := \bar{N}_z \left(\int_0^t \frac{\tilde{\gamma}_z(\xi_s^n)}{\|\tilde{\gamma}_z\|_\infty} ds \right). \quad (3.6)$$

Note that the jump times of this process have the same distribution as N_z^n . Similarly, consider the time-changed process

$$M_z(t) := \bar{N}_z \left(\int_0^t \frac{\tilde{\gamma}_z(\hat{u}(s, 0))}{\|\tilde{\gamma}_z\|_\infty} ds \right), \quad (3.7)$$

and note that its jump times have the same distribution as N_z . Hence, we can assume that there is a coupling under which the jump times of N_z^n and N_z are given by the jump times of (3.6) and (3.7), respectively.

Note now that by definition, τ^n and τ are the first time that any of the processes $\{M_z^n: z \in \mathbb{Z}\}$ and $\{M_z: z \in \mathbb{Z}\}$ have a jump, respectively. Since the time-change in (3.6) converges to the one in (3.7), τ^n converges to τ when conditioning on the realization of $\{\tilde{N}_z: z \in \mathbb{Z}\}$. Moreover, due to (3.4), the jump events of $\{\tilde{N}_z: z \in \mathbb{Z}\}$ are well-separated, consequently, the same holds true for the index z^n of the first jump. Therefore, by the hydrodynamic limit without macroscopic jumps, $x_{[0, \tau^n]}^n$ converges to $x_{[0, \tau]}$, and $x_{\tau^n}^n - x_{\tau^n-}^n = nz^n$ converges to $z := x_\tau - x_{\tau-}$.

It remains to show that the jump rates of x_t match those given in (3.5). Observe that on the event that $\tau \geq t$, for arbitrary $\epsilon > 0$, the probability of the occurrence of a jump of size z before time $t + \epsilon$ is given by

$$\begin{aligned} P(M_z(t + \epsilon) - M_z(t) \geq 1) &= P(M_z(t + \epsilon) - M_z(t) = 1) + o(\epsilon) \\ &= 1 - \exp \int_t^{t+\epsilon} \tilde{\gamma}_z(\hat{u}(s, 0)) \, ds + o(\epsilon) = \int_t^{t+\epsilon} \tilde{\gamma}_z(\hat{u}(s, 0)) \, ds + o(\epsilon). \end{aligned}$$

Moreover, for $s < \tau$, $\hat{u}(s, 0) = u(s, x_s)$. Therefore $x_{[0, \tau^n]}^n$ indeed converges to $x_{[0, \tau]}$ with x_t given by (3.5). \square

References

- [1] L. Avena, F. den Hollander and F. Redig. Large deviation principle for one-dimensional random walk in dynamic random environment: Attractive spin-flips and simple symmetric exclusion. *Markov Process. Related Fields* **16** (2010) 139–168. [MR2664339](#)
- [2] L. Avena, R. dos Santos and F. Völlering. A transient random walk driven by an exclusion process: Regenerations, limit theorems and an Einstein relation. *ALEA Lat. Am. J. Probab. Math. Stat.* **10** (2) (2013) 693–709. [MR3108811](#)
- [3] L. Avena and P. Thomann. Continuity and anomalous fluctuations in random walks in dynamic random environments: Numerics, phase diagrams and conjectures. *J. Stat. Phys.* **147** (2012) 1041–1067. [MR2949519](#)
- [4] F. Comets, N. Gantert and O. Zeitouni. Quenched, annealed and functional large deviations for one-dimensional random walk in random environment. *Probab. Theory Related Fields* **118** (2000) 65–114. [MR1785454](#)
- [5] D. Dolgopyat, G. Keller and C. Liverani. Random walk in Markovian environment. *Ann. Probab.* **36** (2008) 1676–1710. [MR2440920](#)
- [6] A. Greven and F. den Hollander. Large deviations for a random walk in random environment. *Ann. Probab.* **22** (1994) 1381–1428. [MR1303649](#)
- [7] F. Redig and F. Völlering. Random walks in dynamic random environments: A transference principle. *Ann. Probab.* **41** (2013) 3157–3180. [MR3127878](#)
- [8] T. Franco and C. Landim. Hydrodynamic limit of gradient exclusion processes with conductances. *Arch. Ration. Mech. Anal.* **195** (2010) 409–439. [MR2592282](#)
- [9] M. Z. Guo, G. C. Papanicolau and S. R. S. Varadhan. Nonlinear diffusion limit for a system with nearest neighbor interactions. *Comm. Math. Phys.* **118** (1988) 31–59. [MR0954674](#)
- [10] M. Jara, C. Landim and S. Sethuraman. Nonequilibrium fluctuations for a tagged particle in mean-zero one dimensional zero-range processes. *Probab. Theory Related Fields* **145** (2009) 565–590. [MR2529439](#)
- [11] C. Kipnis and C. Landim. *Scaling Limits of Particle Systems. Grundlehren der Mathematischen Wissenschaften* **320**. Springer, Berlin, 1999. [MR1707314](#)
- [12] C. Landim, S. Olla and S. R. S. Varadhan. Asymptotic behavior of a tagged particle in simple exclusion processes. *Bol. Soc. Brasil. Mat.* **31** (3) (2000) 241–275. [MR1817088](#)
- [13] A. S. Sznitman. *Lectures on Random Motions in Random Media. Ten Lectures on Random Media. DMV-Lectures* **32**. Birkhäuser, Basel, 2002. [MR1890289](#)
- [14] O. Zeitouni. Random walks in random environments. *J. Phys. A: Math. Gen.* **39** (2006) 433–464. [MR2261885](#)